

MLL2 and *KDM6A* Mutations in Patients With Kabuki Syndrome

Noriko Miyake,^{1*} Eriko Koshimizu,¹ Nobuhiko Okamoto,² Seiji Mizuno,³ Tsutomu Ogata,⁴ Toshiro Nagai,⁵ Tomoki Kosho,⁶ Hirofumi Ohashi,⁷ Mitsuhiro Kato,⁸ Goro Sasaki,⁹ Hiroyo Mabe,¹⁰ Yoriko Watanabe,¹¹ Makoto Yoshino,¹¹ Toyojiro Matsuishi,¹¹ Jun-ichi Takanashi,¹² Vorasuk Shotelersuk,¹³ Mustafa Tekin,¹⁴ Nobuhiko Ochi,¹⁵ Masaya Kubota,¹⁶ Naoko Ito,¹⁷ Kenji Ihara,¹⁷ Toshiro Hara,¹⁷ Hidefumi Tonoki,¹⁸ Tohru Ohta,¹⁹ Kayoko Saito,²⁰ Mari Matsuo,²⁰ Mari Urano,²⁰ Takashi Enokizono,²¹ Astushi Sato,²² Hiroyuki Tanaka,²³ Atsushi Ogawa,²⁴ Takako Fujita,²⁵ Yoko Hiraki,²⁶ Sachiko Kitanaka,²² Yoichi Matsubara,²⁷ Toshio Makita,²⁸ Masataka Taguri,²⁹ Mitsuko Nakashima,¹ Yoshinori Tsurusaki,¹ Hirotomo Saitsu,¹ Ko-ichiro Yoshiura,³⁰ Naomichi Matsumoto,^{1*} and Norio Niikawa¹⁹

¹Department of Human Genetics, Yokohama City University Graduate School of Medicine, Yokohama, Japan

²Department of Medical Genetics, Osaka Medical Center and Research Institute for Maternal and Child Health, Izumi, Japan

³Department of Pediatrics, Central Hospital, Aichi Human Service Center, Kasugai, Japan

⁴Department of Pediatrics, Hamamatsu University School of Medicine, Hamamatsu, Japan

⁵Department of Pediatrics, Dokkyo Medical University Koshigaya Hospital, Saitama, Japan

⁶Department of Medical Genetics, Shinshu University School of Medicine, Matsumoto, Japan

⁷Division of Medical Genetics, Saitama Children's Medical Center, Iwatsuki, Japan

⁸Department of Pediatrics, Yamagata University Faculty of Medicine, Yamagata, Japan

⁹Department of Pediatrics, Tokyo Dental College Ichikawa General Hospital, Chiba, Japan

¹⁰Department of Child Development, Kumamoto University Hospital, Kumamoto, Japan

¹¹Department of Pediatrics and Child Health, Kurume University School of Medicine, Kurume, Japan

¹²Department of Pediatrics, Kameda Medical Center, Kamogawa, Chiba, Japan

¹³Center of Excellence for Medical Genetics, Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

¹⁴John P. Hussman Institute for Human Genomics, Miller School of Medicine, University of Miami, Miami, Florida

¹⁵Department of Pediatrics, Aichi Prefectural Hospital and Habilitation Center for Disabled Children, Dai-ni Aoitori Gakuen, Okazaki, Japan

¹⁶Division of Neurology, National Center for Child Health and Development, Tokyo, Japan

¹⁷Department of Pediatrics, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

¹⁸Section of Clinical Genetics, Department of Pediatrics, Tenshi Hospital, Sapporo, Japan

¹⁹Research Institute of Personalized Health Sciences, Health Science University of Hokkaido, Tobetsu, Japan

²⁰Institute of Medical Genetics, Tokyo Women's Medical University, Tokyo, Japan

²¹Department of Pediatrics, University of Tsukuba Hospital, Tsukuba, Japan

²²Department of Pediatrics, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

²³Department of Pediatrics, Ohta Nishinouchi Hospital, Tokyo, Japan

²⁴Department of Pediatrics, Chikushi Hospital, Fukuoka University, Fukuoka, Japan

²⁵Department of Pediatrics, Fukuoka University School of Medicine, Fukuoka, Japan

²⁶Hiroshima Municipal Center for Child Health and Development, Hiroshima, Japan

²⁷Department of Medical Genetics, Tohoku University School of Medicine, Sendai, Japan

²⁸Education Center, Asahikawa Medical University, Asahikawa, Japan

²⁹Department of Biostatistics and Epidemiology, Yokohama City University Graduate School of Medicine, Yokohama, Japan

³⁰Department of Human Genetics, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan

Manuscript Received: 9 January 2013; Manuscript Accepted: 9 May 2013

Kabuki syndrome is a congenital anomaly syndrome characterized by developmental delay, intellectual disability, specific facial features including long palpebral fissures and ectropion of the lateral third of the lower eyelids, prominent digit pads, and skeletal and visceral abnormalities. Mutations in *MLL2* and *KDM6A* cause Kabuki syndrome. We screened 81 individuals with Kabuki syndrome for mutations in these genes by conventional methods (n = 58) and/or targeted resequencing (n = 45) or whole exome sequencing (n = 5). We identified a mutation in *MLL2* or *KDM6A* in 50 (61.7%) and 5 (6.2%) cases, respectively. Thirty-five *MLL2* mutations and two *KDM6A* mutations were novel. Non-protein truncating-type *MLL2* mutations were mainly located around functional domains, while truncating-type mutations were scattered through the entire coding region. The facial features of patients in the *MLL2* truncating-type mutation group were typical based on those of the 10 originally reported patients with Kabuki syndrome; those of the other groups were less typical. High arched eyebrows, short fifth finger, and hypotonia in infancy were more frequent in the *MLL2* mutation group than in the *KDM6A* mutation group. Short stature and postnatal growth retardation were observed in all individuals with *KDM6A* mutations, but in only half of the group with *MLL2* mutations. © 2013 Wiley Periodicals, Inc.

Key words: Kabuki syndrome; *MLL2*; *KDM6A*; mutation; genotype–phenotype correlation

INTRODUCTION

Kabuki syndrome (KS; OMIM 147920) is a multiple congenital anomaly syndrome that was originally reported by Niikawa et al. [1981] and Kuroki et al. [1981] (also known as Kabuki make-up syndrome or Niikawa–Kuroki syndrome). KS is diagnosed clinically by characteristic facial features, including long palpebral fissures and ectropion of the lateral third of the lower eyelids, postnatal growth impairment (short stature), developmental delay, intellectual disability, dermatoglyphic abnormalities, visceral and skeletal abnormalities, and immunological dysfunction. The prevalence of the disorder is estimated to be 1 in 32,000 live births [Niikawa et al., 1988]. Two genes have shown to be mutated in patients with KS: *MLL2* (myeloid/lymphoid or mixed-lineage leukemia 2; NM_003482.3) at 12q13.12 and *KDM6A* (lysine (K)-specific demethylase 6A; NM_021140.2) at Xp11.3 [Ng et al., 2010; Lederer et al., 2012; Miyake et al., 2013]. *MLL2* encodes a histone H3 lysine 4 (H3K4)-specific methyl transferase and *KDM6A* is a specific demethylase of histone H3 lysine 27 (H3K27) [Prasad et al., 1997; Lee et al., 2007]. They are both trithorax group proteins and bind each other [Schuettengruber et al., 2007]. These proteins are important for the chromatin state and transcription activation: *MLL2* methylates H3K4 and *KDM6A* removes the H3K27 trimethylation repressive mark

How to Cite this Article:

Miyake N, Koshimizu E, Okamoto N, Mizuno S, Ogata T, Nagai T, Kosho T, Ohashi H, Kato M, Sasaki G, Mabe H, Watanabe Y, Yoshino M, Matsuishi T, Takanashi J-i, Shotelersuk V, Tekin M, Ochi N, Kubota M, Ito N, Ihara K, Hara T, Tonoki H, Ohta T, Saito K, Matsuo M, Urano M, Enokizono T, Sato A, Tanaka H, Ogawa A, Fujita T, Hiraki Y, Kitanaka S, Matsubara Y, Makita T, Taguri M, Nakashima M, Tsurusaki Y, Saitsu H, Yoshiura K-i, Matsumoto N, Niikawa N. 2013. *MLL2* and *KDM6A* mutations in patients with Kabuki syndrome. Am J Med Genet Part A 9999:1–10.

[Dubuc et al., 2013]. The loss of *MLL2* or *KDM6A* function may lead to repressed transcription [Dubuc et al., 2013].

To our knowledge, there has been no comprehensive screen for mutations in these two genes in the same patient series. In this report, we performed a mutation screen of both genes in 81 patients with KS. We then evaluated the clinical features based on the genetic information.

MATERIALS AND METHODS

Samples

Eighty-one individuals clinically suspected to have KS were incorporated in this study: 77 Japanese, two Caucasians, one Belgian, and one Thai. They were all sporadic except for KMS-79, who had an affected sibling. Peripheral blood samples or saliva samples from the

Additional supporting information may be found in the online version of this article at the publisher's web-site.

Conflict of interest: none.

Grant sponsor: Ministry of Health, Labour and Welfare of Japan; Grant sponsor: Japan Science and Technology Agency; Grant sponsor: Strategic Research Program for Brain Sciences; Grant sponsor: Ministry of Education, Culture, Sports, Science and Technology of Japan; Grant sponsor: Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science; Grant sponsor: Grant-in-Aid for Young Scientists from the Japan Society for the Promotion of Science; Grant sponsor: Takeda Science Foundation; Grant sponsor: Yokohama Foundation for the Advancement of Medical Science; Grant sponsor: Hayashi Memorial Foundation for Female Natural Scientists.

*Correspondence to:

Dr. Noriko Miyake or Prof. Naomichi Matsumoto, Department of Human Genetics, Yokohama City University Graduate School of Medicine, 3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan.

E-mails: nmiyake@yokohama-cu.ac.jp (N. Miyake) or naomat@yokohama-cu.ac.jp (N. Matsumoto)

Article first published online in Wiley Online Library (wileyonlinelibrary.com): 00 Month 2013

DOI 10.1002/ajmg.a.36072

patients and their parents (when available) were collected with informed consent and DNA was extracted using a QuickGene-610L (Fujifilm, Tokyo, Japan) or Oragene-DNA kit (DNA Genotek, Inc., Ottawa, Canada) according to the manufacturer's instructions. This study included four previously reported patients (KMS-50, KMS-51, KMS-61, and KMS-71) [Tekin et al., 2006; Torii et al., 2009; Ito et al., 2012]. In addition, three patients with a *KDM6A* mutation were previously described as Patients 1, 2, and 3 by Miyake et al. [2013], and are named KMS-31, KMS-37, and KMS-65, respectively, in this report. This study was approved by the Institutional Review Board of Yokohama City University School of Medicine.

Mutation Screening

Fifty-eight patients (KMS-01 to KMS-69) were screened for *MLL2* mutations by the high-resolution melting (HRM) method using a LightCycler 480 System II (Roche Diagnostics, Indianapolis, IN) and subsequent Sanger sequencing. If an HRM curve pattern was different from those of controls, the DNA sample was Sanger sequenced on an ABI 3500xl or 3130xl Genetic Analyzer (Applied Biosystems, Foster City, CA) and the sequences were analyzed using Sequencher software version 4.10.1 (Gene Codes Corporation, Ann Arbor, MI). *KDM6A* was analyzed in samples with no *MLL2* mutation using HRM analysis and Sanger sequencing as above ($n = 37$). For male samples, genotyping using spike-in control male genomic DNA (10%) was performed to detect a hemizygous mutation. The latter 23 patients (KMS-70 to KMS-92), as well as 22 patients with no mutation in either gene detected by conventional methods, were analyzed by targeted resequencing as described in the following section. We judged a variant as pathogenic when it was previously reported to cause KS, or novel variant when it was not observed in unaffected parents or in in-house exome data ($n = 977$), dbSNP135, or EVS6500 (Exome Variant Server, NHLBI GO Exome Sequencing Project, Seattle, WA; <http://evs.gs.washington.edu/EVS/>; accessed March 1, 2013). In addition, the missense mutation predicted to be polymorphism by both of two predictions (Polyphen-2: <http://genetics.bwh.harvard.edu/pph2/> [Adzhubei et al., 2010] and MutationTaster: <http://www.mutationtaster.org/> [Schwarz et al., 2010]) was considered to be non-pathogenic. Parentage analysis was conducted for the patients only when the parental samples were available. TaKaRa Ex Taq and TaKaRa LA Taq (both Takara, Tokyo, Japan) were used for amplification. The primer sequences and PCR conditions are available on request. All pathological variants were confirmed by Sanger sequencing. Nucleotide numbering reflects cDNA numbering with +1 corresponding to the A of the ATG translation initiation codon in the reference sequence (RefSeq NM_003482.3 for *MLL2*, RefSeq NM_021140.2 for *KDM6A*).

Targeted Resequencing of *MLL2* and *KDM6A* by Next-Generation Sequencing

Ion AmpliSeq Custom Panels (Life Technologies, Inc., Grand Island, NY) covering the entire coding region of *MLL2* and *KDM6A* were created via the Ion AmpliSeq Designer v1.2 (<https://ampliseq.com/browse.action>). Libraries were prepared using the Ion AmpliSeq Library Kit 2.0 (Life Technologies, Inc.), with 10 ng of genomic DNA for each primer pool (two pools for this

analysis). An Agilent 2200 TapeStation (Agilent Technologies, Santa Clara, CA) and the associated High Sensitivity D1K Screen Tape (Agilent Technologies) were used to check the size distribution and the concentration of the DNA libraries. Emulsion PCR and enrichment steps were carried out using the Ion OneTouch 200 Template Kit v2 (Life Technologies, Inc.). The amplicon libraries were sequenced on an Ion Torrent Personal Genome Machine system using 314 or 316 chips, and bar-coding was applied with an Ion Xpress Barcode Adapters 1–16 Kit (all Life Technologies, Inc.). Torrent Suite 2.2 (Life Technologies, Inc.) was used for mapping, base calling, and variant calling. Sequences were annotated using SeattleSeq Annotation 134 (<http://snp.gs.washington.edu/SeattleSeqAnnotation134/>). All variants were confirmed by Sanger sequencing.

Whole Exome Sequencing by High-Throughput Next-Generation Sequencing

Whole exome sequencing was performed in five individuals (KMS-09, -18, -21, -23, and -61) who had no *MLL2* or *KDM6A* abnormality by HRM analysis. DNA was processed with a SureSelect Human All Exon V4 kit (Agilent Technologies), sequenced on a HiSeq2000 (Illumina, Inc., San Diego, CA), and analyzed as previously described [Tsurusaki et al., 2013]. Variants in *MLL2* and *KDM6A* were confirmed by Sanger sequencing.

X-Inactivation Assay

X-inactivation analysis was performed as described [Allen et al., 1992] with slight modification. Briefly, genomic DNA (500 ng) was digested with two methylation-sensitive restriction enzymes, *HpaII* and *HhaI* (New England Biolabs, Beverly, MA), and purified by phenol/chloroform extraction and ethanol precipitation. Digested and undigested DNA samples (10 ng) were separately amplified for the (CAG) n polymorphism at the androgen receptor locus. The forward primer was labeled with 5' FAM dye. PCR products were analyzed on an ABI 3500xl Genetic Analyzer using GeneMapper Software Version 4.1 (Applied Biosystems). The assay was independently performed twice.

cDNA Sequencing

Total RNA was extracted from a lymphoblastoid cell line established from KMS-81 (c.1909_1912del in *KDM6A*) using an RNeasy Plus mini kit (Qiagen, Hilden, Germany) with and without cycloheximide treatment (30 μ g/ml) for 4 hr before cell collection. Reverse transcription (RT) was performed using a Superscript III First-Strand synthesis system for RT-PCR (Life Technologies, Inc.). As the mutation was located in exon 17, the region from the exon 15/16 boundary to the exon 17/18 boundary of *KDM6A* was amplified using cDNA-specific primer pairs (sequences available on request) and sequenced by the Sanger method.

Statistical Analysis

The frequencies of clinical features in the two groups were compared by Fisher's exact test. A difference was considered statistically




FIG. 1. *MLL2* mutations in patients with Kabuki syndrome. Upper: Non-truncating-type mutations. Middle: *MLL2* protein structure with functional domains. The protein contains seven plant homeodomains (PHD), one high-mobility group (HMG) domain, a Phe-Tyr-rich N-terminal (FYRN) domain, a Phe-Tyr-rich C-terminal (FYRC) domain, a SET [Suvar3-9, Enhancer-of-zeste, Trithorax] domain, and a PostSET domain. These functional domains were based on a prediction by SMART (<http://smart.embl-heidelberg.de/>) and the UniProtKB database (<http://www.uniprot.org/uniprot/O14686>). Lower: Truncating-type mutations. Black circles denote nonsense mutations or frameshift mutations. Stars represent splice-site mutations.

significant if $P < 0.05$. Correction for multiple testing was not applied.

RESULTS

Overall Mutation Detection Rates

Pathogenic mutations in *MLL2* and *KDM6A* were found in 50 (61.7%) and five (6.2%) of the 81 patients with KS, respectively (Figs. 1 and 2, Tables I and II). Of the 50 *MLL2* mutations, 35 (70.0%) were predicted to be protein truncating-type and 15 (30.0%) were predicted to be non-truncating-type. Interestingly, non-truncating mutations were mostly localized in or adjacent to the functional domains, while truncating mutations were scattered

throughout the entire coding region (Fig. 1). Fifteen of the *MLL2* mutations have been previously reported (Table I). Three novel variants (not included in the 50 mutations) were considered non-pathogenic (Supplemental Table I). Variant c.10942C > G in patient KMS-22 was predicted to be benign by Polyphen-2 and MutationTaster, c.8813C > T in patient KMS-62 was inherited from an unaffected father, and c.4065A > T in KMS-75 was found heterozygously in our 977 in-house controls. An in-frame duplication in patients KMS-40 and KMS-62, which predicted p.Asp5055_Leu5056dup, was predicted to be polymorphic by MutationTaster, but was previously reported as a pathogenic mutation [Micale et al., 2011]. In addition, the other in-frame mutation in KMS-02 was also predicted to be polymorphic. Unfortunately, parental samples were unavailable for these individuals, except for the mother of patient KMS-62, who did not have this mutation; thus, the de novo status remains unclear. Of the five *KDM6A* mutations including three mutations reported previously [Miyake et al., 2013], four were truncating-type and one was an in-frame deletion located within the Jumonji C domain (Fig. 2).

Clinical Comparison Between the Mutation-Positive and -Negative Groups

We compared the clinical features of the *MLL2* or *KDM6A* mutation-positive and -negative groups (Supplemental Table II). Long palpebral fissures were observed in almost all patients. Cleft lip/palate was more frequently observed in the mutation-positive group ($P = 0.0197$). Interestingly, developmental delay and intellectual disability were observed in all individuals with mutations but were unobserved in some mutation-negative cases ($P = 0.0314$




FIG. 2. *KDM6A* mutations in patients with Kabuki syndrome. Upper: Scheme of *KDM6A* protein. It contains a Jumonji C [JmjC] domain [purple box]. Lower: Black circles indicate truncating-type mutations (nonsense or frameshift mutations); the white circle represents a non-truncating-type mutation (in-frame deletion).

TABLE I. *MLL2* and *KDM6A* Mutations in Patients With KS

Patient ID	Method	Mutation	Predicted amino acid change	De novo	Remarks ^a
Patients with <i>MLL2</i> mutations					
KMS-02	H	c.9831_9848dup	p.Gln3277_Gln3282dup	Unknown	Novel
KMS-08	H	c.12688C > T	p.Gln4230*	Yes	Hannibal et al. [2011]
KMS-13	H	c.2433_2434insCA	p.Glu812Glnfs*119	Yes	Novel
KMS-14	H	c.11806_11807dup	p.Gln3936Hisfs*44	Yes	Novel
KMS-15	H	c.15119A > G	p.Asp5040Gly	Yes	Novel
KMS-17	H	c.5707C > T	p.Arg1903*	Yes	Novel
KMS-18	W	c.12151delA	p.Ile4051*	Yes	Novel
KMS-20	H	c.1300delC	p.Leu434*	Unknown	Novel
KMS-21	W	c.3326_3336dup	p.Asp1113Profs*10	Unknown	Novel
KMS-22	H	c.4127T > G	p.Met1376Arg	Unknown	Novel
KMS-23	W	c.15461G > A	p.Arg5154Gln	Unknown	Li et al. [2011]
KMS-24	H	c.2608 G > T	p.Glu870*	Unknown	Novel
KMS-25	H	c.11917C > T	p.Gln3973*	Unknown	Novel
KMS-27	H	c.15142C > T	p.Arg5048Cys	Yes	Hannibal et al. [2011], Makrythanasis et al. [2013]
KMS-28	H	c.14861C > A	p.Ser4954*	Unknown	Novel
KMS-29	H	c.4419-16 > T	splice site	Unknown	Novel
KMS-30	H	c.4633C > T	p.Gln1545*	Unknown	Novel
KMS-32	H	c.8059C > T	p.Arg2687*	Unknown	Banka et al. [2012b]
KMS-33	T	c.11962C > T	p.Gln3988*	Unknown	Novel
KMS-36	H	c.4736_4737delAG	p.Glu1579Alafs*23	Unknown	Novel
KMS-38	H	c.15143G > A	p.Arg5048His	Unknown	Makrythanasis et al. [2013]
KMS-40	T	c.15163_15168dup	p.Asp5055_Leu5056dup	Unknown	Micale et al. [2011]
KMS-41	H	c.1328delC	p.Pro443Hisfs*487	Yes	Ng et al. [2010]
KMS-42	H	c.16052G > A	p.Arg5351Gln	Yes	Novel
KMS-43	H	c.510 + 1G > A	splice site	Unknown	Novel
KMS-49	T	c.15565G > A	p.Gly5189Arg	Unknown	Novel
KMS-51 ^b	H	c.6297_6298delAC	p.Pro2100Glyfs*54	Yes	Novel
KMS-52	H	c.4693 + 1G > T	splice site	Yes	Novel
KMS-53	H	c.10090C > T	p.Gln3364*	Yes	Novel
KMS-54	T	c.8401C > T	p.Arg2801*	Yes	Novel
KMS-56	H	c.15536G > A	p.Arg5179His	Yes	Ng et al. [2010]
KMS-58	H	c.4333T > G	p.Cys1445Gly	Yes	Novel
KMS-59	H	c.15256C > T	p.Arg5086*	Yes	Banka et al. [2012b]
KMS-60	H	c.11761C > T	p.Gln3921*	Unknown	Novel
KMS-61 ^c	W	c.5269C > T	p.Arg1757*	Yes	Novel
KMS-62	H	c.15163_15168dup	p.Asp5055_Leu5056dup	Unknown ^f	Micale et al. [2011]
KMS-63	T	c.4577G > T	p.Cys1526Phe	Yes	Novel
KMS-69	H	c.11944C > T	p.Arg3982*	Yes	Paulussen et al. [2011]
KMS-70	T	c.13903C > T	p.Gln4635*	Yes	Novel
KMS-71 ^d	T	c.12220C > T	p.Gln4074*	Unknown	Novel
KMS-72	T	c.15061C > T	p.Arg5021*	Yes	Banka et al. [2012b]
KMS-73	T	c.12274C > T	p.Gln4092*	Unknown	Micale et al. [2011]
KMS-76	T	c.4490_4491delAC	p.His1497Leufs*30	Yes	Novel
KMS-78	T	c.16338 + 1G > T	splice site	Yes	Novel
KMS-80	T	c.15088C > T	p.Arg5030Cys	Unknown	Makrythanasis et al. [2013]
KMS-82	T	c.3511G > T	p.Glu1171*	Yes	Novel
KMS-85	T	c.11722C > T	p.Gln3908*	Yes	Paulussen et al. [2011]
KMS-87	T	c.3281_3282delTC	p.Leu1094Profs*20	Yes	Novel
KMS-88	T	c.16052 + 1G > C	splice site	Unknown	Novel
KMS-91	T	c.4267C > T	p.Arg1423Cys	Unknown	Novel
Patients with <i>KDM6A</i> mutations					
KMS-31 ^e	H	c.3717G > A	p.Trp1239*	Unknown	Miyake et al. [2013]
KMS-37 ^e	H	c.1555C > T	p.Arg519*	Unknown	Miyake et al. [2013]
KMS-65 ^e	H	c.3354_3356delTCT	p.Leu1119del	Yes	Miyake et al. [2013]
KMS-81	T	c.1909_1912delTCTA	p.Ser637Thrfs*53	Yes	Novel
KMS-83	T	c.4051C > T	p.Arg1351*	Unknown	Novel

H, high-resolution melting analysis/Sanger sequencing; T, targeted resequencing; W, whole exome sequencing. RefSeq NM_003482.3 for *MLL2* and RefSeq NM_021140.2 for *KDM6A* were used as reference sequences.

^aReferences are listed when the same mutation has been reported previously.

^bThis patient was reported as proband 1 by Tekin et al. [2006].

^cThe detailed clinical features of this patient were reported by Ito et al. [2013] because of her hypothalamic pituitary complications.

^dThe clinical course of this patient, particularly the idiopathic thrombocytopenic purpura, was reported by Torii et al. [2009].

^eThese patients have been reported in our previous study [Miyake et al., 2013].

^fPatient KMS-62: no mutation in the mother.

TABLE II. *MLL2* Non-Truncating-Type Mutations in Patients With KS

Amino acid change ^a	Patient ID	Domain	Polyphen-2 (score)	MutationTaster
p.Met1376Arg	KMS-22	—	Probably damaging (0.915)	Polymorphism
p.Arg1423Cys	KMS-91	PHD	Probably damaging (1.000)	Disease causing
p.Cys1445Gly	KMS-58	PHD	Probably damaging (1.000)	Disease causing
p.Cys1526Phe	KMS-63	PHD	Probably damaging (0.999)	Disease causing
p.Gln3277_Gln3282dup	KMS-02	—	NA	Polymorphism
p.Arg5030Cys	KMS-80	—	Probably damaging (1.000)	Disease causing
p.Asp5040Gly	KMS-15	—	Probably damaging (1.000)	Disease causing
p.Arg5048Cys	KMS-27	—	Probably damaging (1.000)	Disease causing
p.Arg5048His	KMS-38	—	Probably damaging (1.000)	Disease causing
p.Asp5055_Leu5056dup	KMS-40, 62	—	NA	Polymorphism
p.Arg5154Gln	KMS-23	—	Probably damaging (1.000)	Disease causing
p.Arg5179His	KMS-56	FYRN	Possibly damaging (0.840)	Disease causing
p.Gly5189Arg	KMS-49	FYRN	Probably damaging (1.000)	Disease causing
p.Arg5351Gln	KMS-42	—	Probably damaging (1.000)	Disease causing

^aThe nucleotide mutation nomenclature for these predicted protein mutations are included in Table I.

and $P = 0.1778$, respectively). Blue sclera, lower lip pits, spine/rib abnormality, hip joint dislocation, umbilical hernia, kidney dysfunction, cryptorchidism, liver abnormality, spleen abnormality, premature thelarche, neonatal hyperbilirubinemia, and anemia were observed only in the mutation-positive group.

Clinical Comparison of the *MLL2*-Mutated and *KDM6A*-Mutated Groups

We compared the clinical features between the *MLL2*-mutated and *KDM6A*-mutated groups (Figs. 3–5, Supplemental Table III). High arched eyebrows, short fifth fingers, and hypotonia in infancy were more frequent in individuals with *MLL2* mutations than in individuals with *KDM6A* mutations ($P = 0.0364$, 0.0039 , and 0.0283 , respectively). Short stature was more frequent in individuals with *KDM6A* mutations ($P = 0.0485$). Although not statistically significant, postnatal growth retardation was observed in all individuals with *KDM6A* mutations, whereas this was observed in only half of the individuals with *MLL2* mutations.

Clinical Comparison of Individuals With a *MLL2* Truncating-Type and Non-Truncating-Type Mutation

Most clinical features were observed at a similar ratio in both groups (Supplemental Table IV), except for prominent ears and hypotonia, which were more frequently observed in the truncating-type group than in the non-truncating-type group ($P = 0.0339$ and $P = 0.0248$, respectively). However, the facial appearance of individuals in the truncating-type group was more typical, based on the ten originally reported patients with KS [Kuroki et al., 1981; Niikawa et al., 1981], than that in the non-truncating-type group (Figs. 3 and 4). Except for patient KMS-58, the facial appearance of patients with a non-truncating-type mutation was rather less typical. It should be noted that these patients had thick eyebrows (not present in patient KMS-56). Furthermore, ectropion of the

lower eyelid, depressed nasal tip, short columella, and prominent ears all seemed less obvious in the individuals with a non-truncating-type mutation.

X-Inactivation Pattern in Female Patients With a *KDM6A* Mutation

A *KDM6A* mutation was identified in two females (KMS-65 and KMS-81). Individual KMS-65 (c.3354_3356del, which predicts p. Leu1119del) showed a random X-inactivation pattern [Miyake et al., 2013], while individual KMS-81 with a frame-shift mutation showed marked skewing (98:2; Supplemental Fig. 1A). By RT-PCR using mRNA derived from a lymphoblastoid cell line from patient KMS-81, we confirmed that both the mutated and normal alleles were transcribed at similar levels when nonsense-mediated mRNA decay (NMD) was inhibited by cycloheximide treatment (Supplemental Fig. 1B). In untreated cells, or cells treated with dimethylsulfoxide (negative control), the mutant allele was transcribed at a lower level than the wild-type allele, indicating that NMD partially eliminated the mutant.

Exome Sequencing

Among the five patients who were also analyzed by whole exome sequencing, mutations were identified and later confirmed by Sanger sequencing in four (Table I). Fragments with an 11 base-pair insertion (c.3326_3336dup) of *MLL2* in patient KMS-21 could not be amplified by Ex Taq, but could be amplified by LA Taq with LA buffer and confirmed by Sanger sequencing. Three other mutations were missed by HRM analysis (Table I).

DISCUSSION

We identified 50 *MLL2* and five *KDM6A* mutations among 81 patients with KS and add to the 246 *MLL2* mutations described in patients with KS [Ng et al., 2010; Hannibal et al., 2011; Li et al., 2011; Micale et al., 2011; Paulussen et al., 2011; Banka et al., 2012a,b;



FIG. 3. Clinical features of patients with Kabuki syndrome harboring a *MLL2* truncating-type mutation. **A:** Facial features of patients with Kabuki syndrome with a *MLL2* truncating-type mutation. The seven panels for patient KMS-54 show serial images at 0, 1, 4, 6.5, 9.5, 22, and 33 years of age, respectively. **B:** Complete right cleft lip/palate and lower lip pits (arrows) in patient KMS-41. **C:** Abnormal dentition in patients KMS-52 and KMS-59. They also showed hypodontia with wide interdendum. **D:** Patient KMS-72 had congenital strabismus and blepharoptosis as well as opacification of the cornea due to Peters anomaly (arrow). **E:** Hand images show short fifth fingers and prominent digit pads (white arrows).



FIG. 4. Clinical features of patients with Kabuki syndrome with a *MLL2* non-truncating-type mutation. **A:** Facial features of patients with KS and a *MLL2* non-truncating-type mutation. **B:** Patient KMS-56 showed abnormal dentition [hypodontia with wide interdentium].

Kokitsu-Nakata et al., 2012; Tanaka et al., 2012; Bogershausen and Wollnik, 2013; Makrythanasis et al., 2013] (Human Gene Mutation Database Professional 2012.3; <https://portal.biobase-international.com/hgmd/pro/gene.php>). Our mutation-positivity rate for either gene was 67.9% (55/81), and that for *MLL2* only was 61.7% (50/81); these figures are compatible with those reported in a review (55–80%) [Banka et al., 2012b]. Mutation-negative patients suggest the existence of unknown genes to cause KS or misdiagnosis.

As for the phenotype–genotype relationship, Banka et al. [2012b] suggested that feeding problems, kidney anomalies, premature thelarche, joint dislocation, and palatal malformation were more frequently observed in patients with *MLL2*-mutations than in patients with normal *MLL2* sequence. Hannibal et al. [2011] reported that renal anomalies were more common in patients who had *MLL2* mutations compared to those who did not. Li

et al. [2011] reported that short stature and renal anomalies were more frequent in patients with *MLL2*-mutations than in those with normal *MLL2* sequence. In our study, premature thelarche was observed only in patients with *MLL2* mutations, but this was not significant ($P = 0.1137$). The frequencies of kidney anomalies, hip joint dislocation, and short stature were not different when comparing those with and without *MLL2* mutations ($P = 0.3030$, $P = 1.0000$, and $P = 0.0717$, respectively; Supplemental Table V). High arched eyebrows, palatal malformation (cleft palate/lip), low posterior hairline, and short fifth finger were more frequently observed in individuals with *MLL2* mutations than in patients with normal *MLL2* ($P = 0.0118$, $P = 0.0284$, $P = 0.0493$, and $P = 0.0137$, respectively; Supplemental Table V).

X-inactivation skewing in patients with KS has been discussed since the discovery of the *KDM6A* deletion in a female with KS




FIG. 5. Clinical features of patients with Kabuki syndrome with a *KDM6A* truncating-type mutation. **A:** Facial features of patient KMS-81 harboring a *KDM6A* mutation [c.1909_1912del, p.Ser637Thrfs*53]. She showed large front teeth with wide interdentium. **B:** Hand image of KMS-81. Short fifth finger was not remarkable.

[Lederer et al., 2012; Miyake et al., 2013]. In two female patients reported here, patient KMS-65, who had an in-frame deletion, showed a random X-inactivation pattern, but patient KMS-81, who had a truncating-type mutation, showed marked skewing. X-inactivation skewing was also reported in two affected females with *KDM6A* deletion reported by Lederer et al. [2012]. cDNA sequence analysis of patient KMS-81 indicated that the mutant allele of *KDM6A* was expressed at a similar level to the wild-type allele under NMD inhibition. This result suggests that *KDM6A* mostly escapes X-inactivation in female lymphoblastoid cells. Interestingly, *KDM6A/Kdm6a* escapes X-inactivation in humans and mice, and in mice its expression level from the inactive X chromosome (Xi) was reported as 15–35% of that from the active X chromosome (Xa) [Greenfield et al., 1998; Xu et al., 2008]. We calculated the hypothetical expression assuming a 30% *KDM6A* expression level from Xi and 100% expression from Xa (Supplemental Fig. 2). In patient KMS-81, who showed marked skewing (98:2), either the mutant X chromosome or the wild-type X was inactivated in 98% of cells. If the mutant were inactivated, the expression level would be below 1 ($1.0 \times 0.98 + 0.3 \times 0.02 = 0.986$). If the wild-type were inactivated, the expression level would also be below 1 ($1.0 \times 0.02 + 0.3 \times 0.98 = 0.314$). The KS phenotype is usually unassociated with Turner syndrome (45,X), with the *KDM6A* expression level at 1.0 [Miyake et al., 2013]. It is possible that having a *KDM6A* expression level of 1.0 is essential for a normal human phenotype. Similarly, males with only one copy of *KDM6A* do not manifest KS. We previously mentioned the possibility of *UTY* compensation for *KDM6A* (Supplemental Fig. 2) [Miyake et al., 2013], although human *UTY* lacks demethylase activity [Hong et al., 2007; Lan et al., 2007]. The recent evidence that $X^{Utx-}X^{Utx-}$ homozygous mice demonstrated a more severe phenotype than $X^{Utx-}Y^{Uty+}$ mice indicates that *UTY* can compensate for the loss of *UTX* in embryonic development [Shpargel et al., 2012]. Because mouse and human *UTY* show 75% identity, and 95% identity in the Jumonji C domain [Shpargel et al., 2012], it is likely that normal human males who have only one copy of *KDM6A* are supplemented by *UTY* in a demethylase-independent manner.

Interestingly, $X^{Utx-}Y^{Uty+}$ mice showed small body size [Shpargel et al., 2012]. Similarly, the human *KDM6A*-mutated group exhibited short stature and postnatal growth retardation.

Regarding our mutation detection methods, HRM analysis and Sanger sequencing are both imperfect. Next-generation sequencing is more sensitive (especially for single nucleotide variants and small insertions/deletions), faster, and cheaper due to multiple gene screening and the potential to multiplex. However, a microdeletion involving *MLL2* or *KDM6A* or low-level mosaicism of a single nucleotide variant might be missed by this method. Therefore, in patients who test mutation-negative, more comprehensive approaches might be necessary. In conclusion, we investigated *MLL2* and *KDM6A* mutations and their clinical consequences in patients with KS. The majority of the clinical features were observed at a similar frequency among patients with either *MLL2* or *KDM6A*-mutations. The genetic basis of the patients who tested mutation-negative (20–45%) remains elusive. Further studies are necessary to understand the whole picture of the genetic aspects of KS and its genotype–phenotype relationships.

ACKNOWLEDGMENTS

We thank the patients and their parents for participating in this work. We also thank Ms. Y. Yamashita, Ms. E. Koike, Ms. S. Sugimoto, Ms. N. Watanabe, Ms. K. Takabe, and Mr. T. Miyama for their technical assistance. This work was supported by research grants from the Ministry of Health, Labour and Welfare of Japan (H. Saitsu, N. Matsumoto, N. Miyake), the Japan Science and Technology Agency (N. Matsumoto), the Strategic Research Program for Brain Sciences (N. Matsumoto), a Grant-in-Aid for Scientific Research on Innovative Areas-(Transcription cycle)-from the Ministry of Education, Culture, Sports, Science and Technology of Japan (N. Matsumoto), a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (N. Matsumoto), a Grant-in-Aid for Young Scientists from the Japan Society for the Promotion of Science (H.S., N. Miyake), the Takeda Science Foundation (N. Matsumoto, N. Miyake), the Yokohama Foundation for the Advancement of Medical Science (N. Miyake), and the Hayashi Memorial Foundation for Female Natural Scientists (N. Miyake).

REFERENCES

- Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS, Sunyaev SR. 2010. A method and server for predicting damaging missense mutations. *Nat Methods* 7:248–249.
- Allen RC, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW. 1992. Methylation of *HpaII* and *HhaI* sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *Am J Hum Genet* 51:1229–1239.
- Banka S, Howard E, Bunstone S, Chandler K, Kerr B, Lachlan K, McKee S, Mehta S, Tavares A, Tolmie J, Donnai D. 2012a. *MLL2* mosaic mutations and intragenic deletion-duplications in patients with Kabuki syndrome. *Clin Genet* 83:467–471.
- Banka S, Veeramachaneni R, Reardon W, Howard E, Bunstone S, Ragge N, Parker MJ, Crow YJ, Kerr B, Kingston H, Metcalfe K, Chandler K, Magee A, Stewart F, McConnell VP, Donnelly DE, Berland S, Houge G, Morton JE, Oley C, Revencu N, Park SM, Davies SJ, Fry AE, Lynch SA, Gill H, Schweiger S, Lam WW, Tolmie J, Mohammed SN, Hobson E, Smith A, Blyth M, Bennett C, Vasudevan PC, Garcia-Minaur S, Henderson A, Goodship J, Wright MJ, Fisher R, Gibbons R, Price SM, Cds D, Temple IK, Collins AL, Lachlan K, Elmslie F, McEntagart M, Castle B, Clayton-Smith J, Black GC, Donnai D. 2012b. How genetically heterogeneous is Kabuki syndrome?: *MLL2* testing in 116 patients, review and analyses of mutation and phenotypic spectrum. *Eur J Hum Genet* 20:381–388.
- Bogershausen N, Wollnik B. 2013. Unmasking Kabuki syndrome. *Clin Genet* 83:201–211.
- Dubuc AM, Remke M, Korshunov A, Northcott PA, Zhan SH, Mendez-Lago M, Kool M, Jones DT, Unterberger A, Morrissy AS, Shih D, Peacock J, Ramaswamy V, Rolider A, Wang X, Witt H, Hielscher T, Hawkins C, Vibhakar R, Croul S, Rutka JT, Weiss WA, Jones SJ, Eberhart CG, Marra MA, Pfister SM, Taylor MD. 2013. Aberrant patterns of H3K4 and H3K27 histone lysine methylation occur across subgroups in medulloblastoma. *Acta Neuropathol* 125:373–384.
- Greenfield A, Carrel L, Pennisi D, Philippe C, Quaderi N, Siggers P, Steiner K, Tam PP, Monaco AP, Willard HF, Koopman P. 1998. The *UTX* gene escapes X inactivation in mice and humans. *Hum Mol Genet* 7:737–742.
- Hannibal MC, Buckingham KJ, Ng SB, Ming JE, Beck AE, McMillin MJ, Gildersleeve HI, Bigham AW, Tabor HK, Mefford HC, Cook J, Yoshiura K, Matsumoto T, Matsumoto N, Miyake N, Tonoki H, Naritomi K,

- Kaname T, Nagai T, Ohashi H, Kurosawa K, Hou JW, Ohta T, Liang D, Sudo A, Morris CA, Banka S, Black GC, Clayton-Smith J, Nickerson DA, Zackai EH, Shaikh TH, Donnai D, Niikawa N, Shendure J, Bamshad MJ. 2011. Spectrum of MLL2 (ALR) mutations in 110 cases of Kabuki syndrome. *Am J Med Genet Part A* 155A:1511–1516.
- Hong S, Cho YW, Yu LR, Yu H, Veenstra TD, Ge K. 2007. Identification of JmjC domain-containing UTX and JMJD3 as histone H3 lysine 27 demethylases. *Proc Natl Acad Sci USA* 104:18439–18444.
- Ito N, Ihara K, Tsutsumi Y, Miyake N, Matsumoto N, Hara T. 2013. Hypothalamic pituitary complications in Kabuki syndrome. *Pituitary* 16:133–138.
- Kokitsu-Nakata NM, Petrin AL, Heard JP, Vendramini-Pittoli S, Henkle LE, dos Santos DV, Murray JC, Richieri-Costa A. 2012. Analysis of MLL2 gene in the first Brazilian family with Kabuki syndrome. *Am J Med Genet Part A* 158A:2003–2008.
- Kuroki Y, Suzuki Y, Chyo H, Hata A, Matsui I. 1981. A new malformation syndrome of long palpebral fissures, large ears, depressed nasal tip, and skeletal anomalies associated with postnatal dwarfism and mental retardation. *J Pediatr* 99:570–573.
- Lan F, Bayliss PE, Rinn JL, Whetstone JR, Wang JK, Chen S, Iwase S, Alpatov R, Issaeva I, Canaani E, Roberts TM, Chang HY, Shi Y. 2007. A histone H3 lysine 27 demethylase regulates animal posterior development. *Nature* 449:689–694.
- Lederer D, Grisart B, Digilio MC, Benoit V, Crespini M, Ghariani SC, Maystadt I, Dallapiccola B, Verellen-Dumoulin C. 2012. Deletion of KDM6A, a histone demethylase interacting with MLL2, in three patients with Kabuki syndrome. *Am J Hum Genet* 90:119–124.
- Lee MG, Villa R, Trojer P, Norman J, Yan KP, Reinberg D, Di Croce L, Shiekhhattar R. 2007. Demethylation of H3K27 regulates polycomb recruitment and H2A ubiquitination. *Science* 318:447–450.
- Li Y, Bogershausen N, Alanay Y, Simsek Kiper PO, Plume N, Keupp K, Pohl E, Pawlik B, Rachwalski M, Milz E, Thoenes M, Albrecht B, Prott EC, Lehmkuhler M, Demuth S, Utine GE, Boduroglu K, Frankenbusch K, Borck G, Gillissen-Kaesbach G, Yigit G, Wiczorek D, Wollnik B. 2011. A mutation screen in patients with Kabuki syndrome. *Hum Genet* 130:715–724.
- Makrythanasis P, van Bon BW, Stehouwer M, Rodriguez-Santiago B, Simpson M, Dias P, Anderlid BM, Arts P, Bhat M, Augello B, Biamino E, Bongers EM, Del Campo M, Cordeiro I, Cueto-Gonzalez AM, Cusco I, Deshpande C, Frysira E, Louise I, Flores R, Galan E, Gener B, Gilissen C, Granneman SM, Hoyer J, Yntema HG, Kets CM, Koolen DA, Marcelis CL, Medeira A, Micale L, Mohammed S, de Munnik SA, Nordgren A, Reardon SP, Revencu N, Roscioli T, Ruitkamp-Versteeg M, Santos HG, Schoumans J, Schuurs-Hoeijmakers JH, Silengo MC, Toledo L, Vendrell T, van der Burgt I, van Lier B, Zweier C, Reymond A, Trembath RC, Perez-Jurado L, Dupont J, de Vries BB, Brunner HG, Veltman JA, Merla G, Antonarakis SE, Hoischen A. 2013. MLL2 mutation detection in 86 patients with Kabuki syndrome: A genotype–phenotype study. *Clin Genet (In Press)*.
- Micale L, Augello B, Fusco C, Selicorni A, Loviglio MN, Silengo MC, Reymond A, Gumiero B, Zucchetti F, D'Addetta EV, Belligni E, Calcagni A, Digilio MC, Dallapiccola B, Faravelli F, Forzano F, Accadia M, Bonfante A, Clementi M, Daolio C, Douzougou S, Ferrari P, Fischetto R, Garavelli L, Lapi E, Mattina T, Melis D, Patricelli MG, Priolo M, Prontera P, Renieri A, Mencarelli MA, Scarano G, della Monica M, Toschi B, Turolla L, Vancini L, Zatterale A, Gabrielli O, Zelante L, Merla G. 2011. Mutation spectrum of MLL2 in a cohort of Kabuki syndrome patients. *Orphanet J Rare Dis* 6:38.
- Miyake N, Mizuno S, Okamoto N, Ohashi H, Shiina M, Ogata K, Tsurusaki Y, Nakashima M, Saito H, Niikawa N, Matsumoto N. 2013. KDM6A point mutations cause Kabuki syndrome. *Hum Mutat* 34:108–110.
- Ng SB, Bigham AW, Buckingham KJ, Hannibal MC, McMillin MJ, Gildersleeve HI, Beck AE, Tabor HK, Cooper GM, Mefford HC, Lee C, Turner EH, Smith JD, Rieder MJ, Yoshiura K, Matsumoto N, Ohta T, Niikawa N, Nickerson DA, Bamshad MJ, Shendure J. 2010. Exome sequencing identifies MLL2 mutations as a cause of Kabuki syndrome. *Nat Genet* 42:790–793.
- Niikawa N, Matsuura N, Fukushima Y, Ohsawa T, Kajii T. 1981. Kabuki make-up syndrome: A syndrome of mental retardation, unusual facies, large and protruding ears, and postnatal growth deficiency. *J Pediatr* 99:565–569.
- Niikawa N, Kuroki Y, Kajii T, Matsuura N, Ishikiriyama S, Tonoki H, Ishikawa N, Yamada Y, Fujita M, Umemoto H, Iwama Y, Kondoh I, Fukushima Y, Nako Y, Matsui I, Urakami T, Aritaki S, Hara M, Suzuki Y, Chyo H, Sugio Y, Hasegawa T, Yamanaka T, Tsukino R, Yoshida A, Nomoto N, Kawahito S, Aihara R, Toyota S, Ieshima A, Funaki H, Ishitobi K, Ogura S, Furumae T, Yoshino M, Tsuji Y, Kondoh T, Matsumoto T, Abe K, Harada N, Miike T, Ohdo S, Naritomi K, Abushwreb AK, Braun OH, Schmid E. 1988. Kabuki make-up (Niikawa–Kuroki) syndrome: A study of 62 patients. *Am J Med Genet* 31:565–589.
- Paulussen AD, Stegmann AP, Blok MJ, Tserpelis D, Posma-Velter C, Detisch Y, Smeets EE, Wagemans A, Schrandt JJ, van den Boogaard MJ, van der Smagt J, van Haeringen A, Stolte-Dijkstra I, Kerstjens-Frederikse WS, Mancini GM, Wessels MW, Hennekam RC, Vreeburg M, Geraedts J, de Ravel T, Fryns JP, Smeets HJ, Devriendt K, Schrandt Stumpel CT. 2011. MLL2 mutation spectrum in 45 patients with Kabuki syndrome. *Hum Mutat* 32:E2018–E2025.
- Prasad R, Zhadanov AB, Sedkov Y, Bullrich F, Druck T, Rallapalli R, Yano T, Alder H, Croce CM, Huebner K, Mazo A, Canaani E. 1997. Structure and expression pattern of human ALR, a novel gene with strong homology to ALL-1 involved in acute leukemia and to *Drosophila* trithorax. *Oncogene* 15:549–560.
- Schuettengruber B, Chourrout D, Vervoort M, Leblanc B, Cavalli G. 2007. Genome regulation by polycomb and trithorax proteins. *Cell* 128:735–745.
- Schwarz JM, Rodelsperger C, Schuelke M, Seelow D. 2010. MutationTaster evaluates disease-causing potential of sequence alterations. *Nat Methods* 7:575–576.
- Shpargel KB, Sengoku T, Yokoyama S, Magnuson T. 2012. UTX and UTY demonstrate histone demethylase-independent function in mouse embryonic development. *PLoS Genet* 8:e1002964.
- Tanaka R, Takenouchi T, Uchida K, Sato T, Fukushima H, Yoshihashi H, Takahashi T, Tsubota K, Kosaki K. 2012. Congenital corneal staphyloma as a complication of Kabuki syndrome. *Am J Med Genet Part A* 158A:2000–2002.
- Tekin M, Fitoz S, Arici S, Cetinkaya E, Incesulu A. 2006. Niikawa–Kuroki (Kabuki) syndrome with congenital sensorineural deafness: Evidence for a wide spectrum of inner ear abnormalities. *Int J Pediatr Otorhinolaryngol* 70:885–889.
- Torii Y, Yagasaki H, Tanaka H, Mizuno S, Nishio N, Muramatsu H, Hama A, Takahashi Y, Kojima S. 2009. Successful treatment with rituximab of refractory idiopathic thrombocytopenic purpura in a patient with Kabuki syndrome. *Int J Hematol* 90:174–176.
- Tsurusaki Y, Kobayashi Y, Hisano M, Ito S, Doi H, Nakashima M, Saito H, Matsumoto N, Miyake N. 2013. The diagnostic utility of exome sequencing in Joubert syndrome and related disorders. *J Hum Genet* 58:113–115.
- Xu J, Deng X, Watkins R, Distechi CM. 2008. Sex-specific differences in expression of histone demethylases Utx and Uty in mouse brain and neurons. *J Neurosci* 28:4521–4527.